

Gravitoelectromagnetism and other decompositions of the Riemann tensor

Hans-Jürgen Schmidt, Institut für Mathematik, Universität Potsdam

Am Neuen Palais 10, D-14469 Potsdam, Germany

<http://www.physik.fu-berlin.de/~hjschmi>, e-mail: hjschmi@rz.uni-potsdam.de

Abstract

We present gravitoelectromagnetism and other decompositions of the Riemann tensor from the differential-geometrical point of view.

The word gravitomagnetism is now in common use, even if some few people (see e.g. page 1306 of Ref. [1]) proposed to write gravimagnetism instead. Three of the many recent papers on this topic are Refs. [2], [3] and [4]. What is the idea behind this topic ?

The Maxwell field can be represented by the tensor F_{ij} , possessing $4^2 = 16$ real components. By use of antisymmetry, this figure reduces to 6, but even a six-dimensional space is hard to imagine. However, by decomposing it into two 3-vectors – the electric and the magnetic 3-vector – the geometric meaning of all the components can be made clear, and this is the form where the contact with experimentalists becomes possible. Mathematically, these 3-vectors can be obtained by the additional introduction of a unit time-like vector field u^i and well-known transvections with it and with the pseudo-tensor ϵ_{ijkl} .

Now, the analogous procedure shall be made with the gravitational field. We have the Riemann tensor R_{ijkl} , possessing $4^4 = 256$ real components. By use of the known symmetries, this figure reduces to 20, but this twenty-dimensional space is even harder to imagine. In the next pages, I will present five different possibilities how to arrange this set of components to get a better understandable system (for details see e.g. Ref. [5]).

The Riemann tensor R_{ijkl} of a space-time of dimension $n \geq 3$ can be decomposed according to several different criteria:

1. The usual one into the Weyl tensor C_{ijkl} plus a term containing the Ricci tensor R_{ij} plus a term containing the Riemann curvature scalar R .
2. Two trace-less parts plus the trace.
3. The Weyl tensor plus only one additional term.
4. Two divergence-free parts plus the trace.
5. The gravitoelectromagnetic point of view.

One should know, that the mathematical construction of the gravitoelectric field is much elder than the word “gravitoelectric”: Einstein calculated the gravitoelectric correction to the Newtonian potential of the sun - even if Einstein himself never had used that word. Concerning the invariant characterization of gravitomagnetism one should be careful, because non-flat space-times exist, whose curvature invariants are all vanishing, see e.g. Ref. [6].

We use the following two properties of the Riemann tensor

$$R_{ijkl} = -R_{ijlk} \quad (1)$$

$$R_{ijkl} = R_{klij} . \quad (2)$$

The Ricci tensor is the trace of the Riemann tensor:

$$R_{ij} = g^{kl} R_{ikjl} , \quad (3)$$

where g_{kl} denotes the metric of the space-time, and the Riemann curvature scalar is the trace of the Ricci tensor

$$R = g^{kl} R_{kl} . \quad (4)$$

The sign conventions are defined such that in Euclidean signature, the curvature scalar of the standard sphere is positive.

For any symmetric tensor H_{ij} we define another tensor H_{ijkl}^* via

$$H_{ijkl}^* = H_{ik}g_{jl} + H_{jl}g_{ik} - H_{il}g_{jk} - H_{jk}g_{il} . \quad (5)$$

Then the tensor H_{ijkl}^* automatically fulfils the identities eqs. (1) and (2). For the special case $H_{ij} = g_{ij}$ we get the simplified form

$$g_{ijkl}^* = 2g_{ik}g_{jl} - 2g_{il}g_{jk} . \quad (6)$$

1 The usual decomposition

The Weyl tensor C_{ijkl} is the trace-less part of the Riemann tensor, i.e.

$$g^{ik} C_{ijkl} = 0; \quad (7)$$

it vanishes identically for $n = 3$. Using the notation of eqs. (5) and (6) we make the ansatz

$$R_{ijkl} = C_{ijkl} + \alpha R_{ijkl}^* + \beta R g_{ijkl}^*. \quad (8)$$

Then the coefficients α and β have to be specified such that the condition eq. (7) becomes an identity. This condition determines the coefficients α and β uniquely, and the result is the following:

$$\alpha = \frac{1}{n-2} \quad \text{and} \quad \beta = \frac{-1}{2(n-1)(n-2)}. \quad (9)$$

Thus, we get the usual formula

$$R_{ijkl} = C_{ijkl} + \frac{1}{n-2} (R_{ik}g_{jl} + R_{jl}g_{ik} - R_{il}g_{jk} - R_{jk}g_{il}) - \frac{1}{(n-1)(n-2)} R (g_{ik}g_{jl} - g_{il}g_{jk}). \quad (10)$$

2 The decomposition using trace-less parts

In distinction to the previous subsection, we now perform a more consequent decomposition into trace and trace-less parts. To this end we define S_{ij} as the trace-less part of the Ricci tensor, i.e. $g^{ij}S_{ij} = 0$ with $S_{ij} = R_{ij} + \kappa R g_{ij}$ possessing the unique solution $\kappa = -1/n$, i.e.,

$$S_{ij} = R_{ij} - \frac{1}{n} R g_{ij}. \quad (11)$$

Then the analogous equation to eq. (8) is

$$R_{ijkl} = C_{ijkl} + \gamma S_{ijkl}^* + \delta R g_{ijkl}^*. \quad (12)$$

Again, eq. (5) has been applied. Eq. (12) becomes a correct identity if and only if

$$\gamma = \frac{1}{n-2} \quad \text{and} \quad \delta = \frac{1}{2n(n-1)}. \quad (13)$$

So we get

$$R_{ijkl} = C_{ijkl} + \frac{1}{n-2} (S_{ik}g_{jl} + S_{jl}g_{ik} - S_{il}g_{jk} - S_{jk}g_{il}) + \frac{1}{n(n-1)} R (g_{ik}g_{jl} - g_{il}g_{jk}) . \quad (14)$$

3 Decomposition into two parts

Let us define a tensor

$$L_{ij} = R_{ij} + \zeta R g_{ij} \quad (15)$$

such that a parameter ε exists which makes

$$R_{ijkl} = C_{ijkl} + \varepsilon L_{ijkl}^* \quad (16)$$

becoming a true identity. It turns out that this is possible iff

$$\zeta = \frac{-1}{2(n-1)} \quad \text{and} \quad \varepsilon = \frac{1}{n-2} . \quad (17)$$

Thus, we can write eq. (15) as

$$L_{ij} = R_{ij} - \frac{1}{2(n-1)} R g_{ij} \quad (18)$$

and eq. (16) as

$$R_{ijkl} = C_{ijkl} + \frac{1}{n-2} (L_{ik}g_{jl} + L_{jl}g_{ik} - L_{il}g_{jk} - L_{jk}g_{il}) . \quad (19)$$

4 Decomposition into divergence-free parts

Now, besides the identities eqs. (1) and (2), we also use identities involving the covariant derivatives, denoted by a semicolon, of the Riemann tensor. The Bianchi identity reads

$$R_{ijkl;m} + R_{ijlm;k} + R_{ijmk;l} = 0 . \quad (20)$$

Its trace can be obtained by transvection with g^{ik} and reads

$$R_{jl;m} + R^i_{jlm;i} - R_{jm;l} = 0 . \quad (21)$$

It should be mentioned, that the transvection with respect to other pairs of indices does not lead to further identities. The Einstein E_{ij} tensor is defined as

$$E_{ij} = R_{ij} + \lambda R g_{ij} , \quad (22)$$

where λ has to be chosen such that the Einstein tensor is divergence-free, i.e.,

$$E^i_{j;i} = 0 . \quad (23)$$

Using the trace of eq. (21) (again, there is essentially only one such trace),

$$2R^i_{l;i} - R_{;l} = 0 , \quad (24)$$

we uniquely get $\lambda = -1/2$, i.e. the Einstein tensor is

$$E_{ij} = R_{ij} - \frac{1}{2}R g_{ij} . \quad (25)$$

With the ansatz

$$R_{ijkl} = W_{ijkl} + \eta E^*_{ijkl} + \vartheta R g^*_{ijkl} \quad (26)$$

it holds: The coefficients η and ϑ are uniquely determined by the requirements that eq. (26) is an identity, and the divergence of the tensor W_{ijkl} vanishes:

$$W^i_{jkl;i} = 0 . \quad (27)$$

We get the following values of the constants:

$$\eta = 1 \quad \text{and} \quad \vartheta = \frac{1}{4} . \quad (28)$$

and then:

$$\begin{aligned} R_{ijkl} = & W_{ijkl} + E_{ik}g_{jl} + E_{jl}g_{ik} - E_{il}g_{jk} - E_{jk}g_{il} \\ & + \frac{1}{2}R(g_{ik}g_{jl} - g_{il}g_{jk}) \end{aligned} \quad (29)$$

defines a decomposition of the Riemann curvature tensor into the divergence-free tensors W_{ijkl} , E_{ij} , g_{ij} and the scalar R .

It should be mentioned, that for every $n > 2$, the four tensors R_{ij} eq. (3), S_{ij} eq. (11), L_{ij} eq. (18) and E_{ij} eq. (25) represent four different tensors.

It is a remarkable fact, that the coefficients in eqs. (25) and (29) do not depend on the dimension n .

5 The gravitoelectromagnetic point of view

The gravitoelectromagnetic point of view can be obtained by the additional introduction of a unit time-like vector field u^i and well-known transvections with it and with the pseudo-tensor ϵ_{ijkl} .

References

- [1] H. Urbandtke, Gen. Rel. Grav. **35** (2003) 1305
- [2] V. Faraoni, R. Dumse, Gen. Rel. Grav. **31** (1999) 91
- [3] Jian Qi Shen, Gen. Rel. Grav. **34** (2002) 1423
- [4] R. Ruffini, C. Sigismondi (Eds.): Nonlinear gravitodynamics - The Lense-Thirring effect, WSPC Singapore (2003)
- [5] J. Schouten: Ricci-Calculus, Springer-Verlag Berlin (1954)
- [6] H.-J. Schmidt: Consequences of the noncompactness of the Lorentz group, gr-qc/9512007; Int. J. Theor. Phys. **37** (1998) 691